

Association of acute thrombocytopenia with anaphylaxis

Vihitha Thota, MD^a, Sudheer Konduru, MD^a, Sana Mulla, MD^a, Samia Hossain, MD^a, Manaswitha Thota, BS^b, and Rajesh Thirumaran, MD^c

^aDepartment of Internal Medicine, Mercy Catholic Medical Center, Darby, Pennsylvania; ^bSchool of Medicine, Virginia Commonwealth University, Richmond, Virginia; ^cDepartment of Hematology and Oncology, Mercy Catholic Medical Center, Darby, Pennsylvania

ABSTRACT

A commonly seen phenomenon in the hospital and critical care setting is anaphylaxis. This acute systemic inflammatory reaction can lead to anaphylactic shock in severe cases and potentially be fatal. The role of platelets in anaphylactic reactions is not well established; however, platelets, among other mediators such as platelet-activating factor, have been shown to promote a pro-thrombotic state shortly after an acute hypersensitivity reaction. In addition, the aggregation of platelets promoted by platelet-activating factor and other mediators can also lead to thrombocytopenia. We present a case of a 57-year-old woman who developed severe anaphylaxis while receiving chemotherapy with paclitaxel suspended in Cremophor, a well-known allergen. She was profoundly thrombocytopenic following the reaction and was treated with therapeutic anticoagulation, with no thrombus formation.

KEYWORDS Anaphylaxis; Cremophor; paclitaxel; thrombocytopenia; thrombosis

Anaphylaxis is an acute, systemic, potentially life-threatening reaction triggered by many agents, including allergens. While predominantly mast cells contribute to the inflammatory response, other inflammatory modulators such as platelets and platelet-activating factor have also been implicated. Some of these mediators are also involved in the coagulation cascade; therefore, their role can lead to an increased risk of thrombocytopenia and thrombus formation in patients with anaphylaxis.¹ However, this association remains underrecognized. Anaphylaxis is a well-known side effect of Cremophor, a vehicle used to administer paclitaxel.² We present a case of severe anaphylaxis during paclitaxel administration, with subsequent severe thrombocytopenia.

CASE PRESENTATION

A 57-year-old woman with a history of newly diagnosed stage IV lung adenocarcinoma presented to our oncology clinic for her second chemoimmunotherapy session with pembrolizumab, carboplatin, and paclitaxel. Within 6 minutes of starting the paclitaxel infusion, she complained of sudden-onset dyspnea, nausea, and a feeling of her “throat closing in.” At this time, her blood pressure was 141/119 mm Hg; heart rate, 130 to 140 beats/min; respiratory rate,

28 breaths/min; oxygen saturation, 79% on room air; and temperature, 97.1°F. Immediately, we stopped the infusion, started supplemental oxygen, and gave intravenous dexamethasone, methylprednisolone, and intramuscular epinephrine. The patient was taken to the emergency room due to continued respiratory distress and concern for anaphylaxis.

Given her increased work of breathing, she was placed on noninvasive positive pressure ventilation and eventually deescalated to nasal cannula once her respiratory status improved. Vital signs had significantly improved by this time, and laboratory workup was notable for a platelet count of 39 10⁹/L, with repeat bloodwork confirming a platelet count of 45 10⁹/L. Five hours before this encounter, at our infusion clinic, her platelet count was 332 10⁹/L (*Table 1*). Coagulation studies, a comprehensive metabolic panel, and a hemolysis panel were unremarkable. Examination revealed a petechial rash on her bilateral arms extending up to the shoulders.

Hematology noted that the thrombocytopenia was likely secondary to the acute anaphylactic reaction and recommended monitoring it clinically. The patient was on therapeutic enoxaparin for a known pulmonary vein thrombus; the hematologist recommended continuing the anticoagulation since anaphylaxis induces a prothrombotic state. Thus, we continued the enoxaparin, and the platelet count improved to 57 10⁹/L 12 hours after admission (*Table 1*).

Corresponding author: Vihitha Thota, MD, 1500 Lansdowne Ave., Box #40, Darby, PA 19023 (e-mail: vihitha.thota@mercyhealth.org)

The authors report no conflicts of interest. Consent was obtained from the patient for her anonymized information to be published in this article.

Received April 20, 2021; Revised May 14, 2021; Accepted May 17, 2021.

Table 1. Complete blood count before and after the anaphylactic reaction displaying the acute drop and recovery of platelet counts

| Test | Day and time | | | | | |
|--------------------------------|--------------|-----------|-----------|-----------|-----------|------------|
| | 1 at 0900 | 1 at 1400 | 1 at 1700 | 2 at 0600 | 5 at 0900 | 12 at 0900 |
| White blood cells ($10^9/L$) | 7.5 | 21.3 | 20.4 | 18.9 | 11 | 6.8 |
| Red blood cells (g/dL) | 3.14 | 3.51 | 3.25 | 3.00 | 3.03 | 2.83 |
| Hemoglobin (g/dL) | 9.5 | 10.4 | 9.7 | 9.1 | 9.1 | 9.8 |
| Hematocrit (%) | 26.8 | 30.3 | 28 | 26.1 | 26.3 | 28.9 |
| Mean corpuscular volume (fL) | 85.3 | 86.4 | 86.1 | 86.9 | 86.6 | 87.6 |
| Platelet count ($10^9/L$) | 332 | 39 | 45 | 57 | 154 | 406 |
| Mean platelet volume (fL) | 9.1 | 7.1 | 6.6 | 8.3 | 8.8 | 7.9 |
| Neutrophils (%) | 60.5 | 94.2 | 94.9 | 92.2 | 65.5 | 61.2 |
| Lymphocytes (%) | 27.7 | 4.8 | 3.7 | 6.3 | 23.5 | 26.3 |

Oxygen was also weaned at this point, and the patient was discharged with close follow-up. Three days after discharge, her platelet count improved to $154 \times 10^9/L$ (Table 1), with resolution of the petechiae. As anaphylaxis occurred during infusion of paclitaxel suspended in Cremophor, the oncologist changed the regimen to Abraxane, which consists of the same drug bound to albumin. The patient has tolerated this new medication well and, so far, has shown a good response to therapy.

DISCUSSION

Anaphylaxis is an acute, potentially lethal systemic process resulting from an inflammatory reaction to medications, among other agents.³ The mechanism behind most cases of anaphylaxis is immunoglobulin E-mediated.⁴ In anaphylaxis, platelets, among other cells, are activated and release platelet-activating factor. Platelet-activating factor causes platelet aggregation and further release of potent vasoconstrictors in the inflammatory response,⁵ causing increased vascular permeability, circulatory collapse, and decreased cardiac output.⁶ Platelets, once activated, are in a procoagulant state by binding to collagen through glycoprotein VI. Platelets also contribute to the activation of coagulation by providing binding sites for prothrombin and factor XI.²

Clinical manifestations of anaphylaxis include rash, dyspnea, chest pain, tachycardia, nausea, vomiting, urticaria, and angioedema.⁷ Thrombocytopenia has been seen in rare cases of anaphylaxis, mainly when it involves shock.¹ Our patient experienced anaphylaxis during an infusion of Cremophor-containing paclitaxel; it is recognized that the anaphylaxis is due to Cremophor as opposed to the paclitaxel,⁸ mediated by primarily a type 1 hypersensitivity reaction.⁹ A new formulation of paclitaxel, nab-paclitaxel or Abraxane, consists of albumin-bound paclitaxel, eliminating the need for Cremophor. Several trials have shown that

patients with prior anaphylaxis to paclitaxel suspended in Cremophor are able to tolerate Abraxane.¹⁰

According to our research, only three previously reported cases^{1,11} have described the association between anaphylaxis, thrombocytopenia, and thrombus formation. All three cases described a significant decrease in platelets (8%–49%, $P < 0.0001$) and thrombus formation in patients with anaphylaxis, without shock. Unlike the previously described cases, our patient received anticoagulation and did not develop a thrombotic event. Of all the cases, ours also had the most severe drop in platelet count, from 332 to $45 \times 10^9/L$ (87%), indicating that anticoagulation may have been protective against thrombus formation. Therefore, clinicians should consider anticoagulation in patients with anaphylaxis-induced thrombocytopenia.

1. Peppers BP, Vatsayan A, Dalal J, et al. A case series: association of anaphylaxis with a significant decrease in platelet levels and possible secondary risk of thrombosis. *Immun Inflamm Dis*. 2018;6(3): 377–381. doi:10.1002/iid3.224.
2. Heemskerk JW, Bevers EM, Lindhout T. Platelet activation and blood coagulation. *Thromb Haemost*. 2002;88(2):186–193.
3. Brown SGA, Kemp SF, Lieberman PL. Anaphylaxis. In: *Middleton's Allergy: Principles and Practice*. 8th ed. Philadelphia: Elsevier; 2014: 1237–1240.
4. Johansson SG, Bieber T, Dahl R, et al. Revised nomenclature for allergy for global use: report of the nomenclature review committee of the World Allergy Organization, October 2003. *J Allergy Clin Immunol*. 2004;113(5):832–836. doi:10.1016/j.jaci.2003.12.591.
5. Finkelman FD. Anaphylaxis: lessons from mouse models. *J Allergy Clin Immunol*. 2007;120(3):506–517. doi:10.1016/j.jaci.2007.07.033.
6. Gill P, Jindal NL, Jagdis A, et al. Platelets in the immune response: revisiting platelet-activating factor in anaphylaxis. *J Allergy Clin Immunol*. 2015;135(6):1424–1432. doi:10.1016/j.jaci.2015.04.019.
7. Irizarry LD, Luu TH, McKoy JM, et al. Cremophor EL-containing paclitaxel-induced anaphylaxis: a call to action. *Commun Oncol*. 2009; 6(3):132–134. doi:10.1016/S1548-5315(11)70224-8.

8. Picard M, Castells MC. Re-visiting hypersensitivity reactions to taxanes: a comprehensive review. *Clin Rev Allergy Immunol*. 2015;49(2): 177–191. doi:[10.1007/s12016-014-8416-0](https://doi.org/10.1007/s12016-014-8416-0).
9. Volcheck GW, Van Dellen RG. Anaphylaxis to intravenous cyclosporine and tolerance to oral cyclosporine: case report and review. *Ann Allergy Asthma Immunol*. 1998;80(2):159–163. doi:[10.1016/S1081-1206\(10\)62949-3](https://doi.org/10.1016/S1081-1206(10)62949-3).
10. Boulanger J, Boursiquot JN, Cournoyer G, et al. Management of hypersensitivity to platinum- and taxane-based chemotherapy: CEPO review and clinical recommendations. *Curr Oncol*. 2014;21(4): e630–e641. doi:[10.3747/co.21.1966](https://doi.org/10.3747/co.21.1966).
11. Johnson JA, Frith JA, Casselman J, et al. Thromboembolic stroke: a sequela of Hymenoptera venom-induced anaphylaxis. *Ann Allergy Asthma Immunol*. 2016;116(3):262–264. doi:[10.1016/j.anai.2016.01.007](https://doi.org/10.1016/j.anai.2016.01.007).

Avocations



Red-naped sapsucker. Shot with a Canon EOS 5D Mark II, 1/500, F 2.8, ISO 200. Photo copyright © William L. Rayburn, MD (William Rayburn@BSWHealth.org), an obstetrician/gynecologist and chief medical officer of the College Station Region for Baylor Scott & White Health. Dr. Rayburn also serves as chair of the Quality Committee for Baylor Scott & White Holdings.